

# *Uso de herbívoros na gestão da vegetação*



**Antonio Rigueiro Rodríguez**  
*Departamento de Producción Vexetal  
e Proxectos de Enxeñaría*  
*Escola Politécnica Superior de Enxeñaría de Lugo*



***ULEX***  
***9% PB***  
***40% MSD***





*Erica australis*  
6% PB  
16% MSD



*Erica arborea*  
8% PB  
24% MSD


















# PIÑEIRAIS-EUCALIPTAIS

## TIPO DE GANDO/CARGA GANADEIRA

- **cabalos-cabras (pasto leñoso)**
- **vacas-ovellas (pasto herbáceo)**
- **1-4 cabras/ovellas por ha**
- **0,25-0,5 cabalos/vacas por ha**













*MONTE MARCO DA CURRA  
MONFERO  
A CORUÑA*



# **AXUSTE DA CARGA GANDEIRA**

## **animais por ha**

	<b>CARGA XERAL</b>	<b>CARGA INSTANTÁNEA</b>
<b>Inicial</b>	<b>2 cabras</b>	<b>8 cabras</b>
<b>Un ano</b>	<b>2 cabras 1 ovella</b>	<b>8 cabras 4 ovellas</b>
<b>Dous anos</b>	<b>1 cabra 2 ovellas</b>	<b>4 cabras 8 ovellas</b>
<b>Tres anos</b>	<b>1 cabra 3 ovellas</b>	<b>4 cabras 12 ovellas</b>







# **EFECTO DO PASTOREO SOBRE A BIOMASA E ALTURA DO SOTOBOSQUE**

<b>TRATAMENTOS</b>	<b>ALTURA MEDIA SOTOBOSQUE cm</b>	<b>BIOMASA SOTOBOSQUE t ms ha<sup>-1</sup></b>
<b>Antes pastoreo</b>	<b>200</b>	<b>20-30</b>
<b>3 anos de pastoreo</b>	<b>10-15</b>	<b>0,5-2</b>







***MONTE DE SAN BREIXO***  
***PARGA, GUITIRIZ***  
***LUGO***





*Ulex gallii*  
*Ulex europaeus*
















- PASTO RESIDUAL media
- PASTO RESIDUAL continuo
- PASTO RESIDUAL rotacional





***MOITO OBRIGADO***